



Approaches of bisphosphonates and osseointegrated implants: a review of the major outcomes

Igor Luis Rodrigues Volpi^{1,*}, Giovanna Charlo Bezerra¹, João Victor Bartolomei¹, José Augusto Parola da Cruz¹

¹ UNORTE - University Center of Northern São Paulo - Department of Dentistry, Sao Jose do Rio Preto, Sao Paulo, Brazil.

*Corresponding author: Igor Luis Rodrigues Volpi.

UNORTE - University Center of Northern São Paulo - Department of Dentistry, Sao Jose do Rio Preto, Sao Paulo, Brazil.

Email: iigorvolpionto@outlook.com

DOI: <https://doi.org/10.54448/mdnt26S309>

Received: 06-22-2026; Revised: 09-14-2026; Accepted: 09-29-2026; Published: 10-01-2026; MedNEXT-id: e26S309

Editor: Dr. Idiberto José Zotarelli-Filho, MSc, Ph.D., Post-Doctoral.

Abstract

The most common bone disorder found by implant dentists is osteoporosis, a systemic skeletal disorder associated with aging characterized by loss of bone mass, making bones fragile and more susceptible to fractures. The World Health Organisation has defined osteoporosis as a bone mineral density level greater than 2.5 standard deviations below the mean of young normal women. After 60 years of age, a third of the population has this disorder; it occurs twice as often in women as in men. It is estimated that 1.3 million fractures and 133,000 hip fractures occur every year as a result of osteoporosis. The aim of this study was to discuss aspects of the pharmacological action of Bisphosphonates (BP) and their influence on bone tissue when associated with treatment with dental implants. The PRISMA rules were followed. The search was conducted from February to April 2026 across the PubMed, Scopus, Web of Science, Embase, Cochrane, Ovid, Periodicos.com, Harvard Library, and Google Scholar. 64 articles were found, 35 articles were evaluated in full, and 26 were included and developed in the present systematic review study. There are several types of treatments that prevent or prevent the progression of osteoporosis. So, BP, such as alendronate, are inhibitors of bone resorption. Act as controlling the development of osteoporosis by increasing the process of bone density and decreasing its reabsorption, often supporting the process of osseointegration of dental implants.

Keywords: Bisphosphonate. Osseointegration. Dental implant. Osteoporosis. Complications.

Introduction

The aim of Modern Dentistry is to restore normal comfort, function, aesthetics, phonation, and health to the patient. What makes Implantology unique is the ability to achieve this goal; however, the more teeth a patient loses, the more challenging the task becomes [1]. It is estimated that the number of dental implants used in the United States increased by more than 10 times from 1983 to 2002 and another five times from 2000 to 2005 [2]. More than one million dental implants are performed each year [3]. The high need and use of treatments related to implants result from the combined effect of several factors, and the most important are: (1) an aging population with longer life expectancy, and (2) age-related dental loss [3].

In this sense, one of the major causes of osteopenia in women over 60 years of age is estrogen deficiency. This deficiency, associated with aging, causes an osteoporotic picture. Hormone replacement is necessary for adequate treatment of the symptoms of menopause and to prevent possible osteoporosis [2,3]. There are some drugs that help in the treatment of postmenopausal osteoporosis; they are calcitonin, bisphosphonates (BP), and the selective modulators of estrogen receptors [3]. Thus, BP has been the best drug associated with significant improvement in the quality of life of patients with bone diseases such as Paget's disease, bone metastases, osteogenesis imperfecta, hypercalcemia, and even severe osteoporosis [3].

These drugs are used worldwide in cancer patients and are given intravenously as zoledronic acid (Zometa®). They can also be administered orally, such as alendronate (Fosamax®) and risedronate (Actonel®)

for the treatment of postmenopausal osteoporosis [Duarte, Nociti, (2004)]. In 2003, a side effect associated with the use of BP with oral manifestation called Osteonecrosis Associated with BP was described for the first time [4].

In this context, osteoporosis is a prevalent global bone disease in human aging. BPs are commonly used as therapy because they influence calcium metabolism of hard and soft tissues. Mucosal and dermal ulceration with exposure of the underlying bone results from incomplete epithelial recovery due to reduced desmosome formation due to a lack of available calcium. However, pathological situations, such as BP-related jaw osteonecrosis, have been described. This hypothesis states that other situations that require intact functional desmosomes, such as skin healing over chronic pressure points leading to pressure ulcers and hemidesmosomes such as epithelial seals in contact with titanium surfaces, will have a higher prevalence of collapse among patients treated with BP. This can be proven by the decreased modulation of calcium ions due to BP and its effect on intercellular communicating junction formation [5-11].

As yet another example of literary support, one paper reported a type of localized osteonecrosis that can occur in patients who had successfully osseointegrated implants for many years and then started anti-resorptive therapy [12]. Eleven female patients who successfully had implants placed but were placed on anti-resorptive therapy (BP or denosumab) several years later and developed osteonecrosis around the implants were identified. In each case, osteonecrosis occurred only around the implants and not around the patient's remaining teeth. Implants from eight patients were removed with bone sequestration firmly attached to the implant. This is different from the normal pattern of implant failure. Implant failure can occur when patients with successfully integrated implants are subsequently placed on anti-resorption therapy, and osteonecrosis takes on a particular form where a sequestration forms that remains adherent to the implant. Why the remaining adjacent teeth are not affected is unclear [2].

Therefore, the present study reviewed the literature analyzing the use of BP associated with dentistry. Because it is a possible alternative for the reduction of loss and increase in bone density, as well as for the possibility of osteonecrosis, which requires therapeutic and preventive measures in the involvement of invasive practices such as dental implants.

Methods

Experimental and clinical studies were included (case reports, retrospective, prospective, and

randomized trials) with qualitative and / or quantitative analysis. Initially, the keywords were determined by searching the DeCS tool (Descriptors in Health Sciences, BIREME base) and later verified and validated by the MeSH system (Medical Subject Headings, the US National Library of Medicine) in order to achieve consistent search results.

Mesh Terms (Descriptors)

The descriptors included were *Bisphosphonate*. *Osseointegration*. *Dental implant*. *Osteoporosis*. *Complications*. The literature search was conducted through online databases: PubMed, Scopus, Web of Science, Embase, Cochrane, Ovid, Periodicos.com, Harvard Library, and Google Scholar. The risk of BIAS in the present study was not analyzed. The PRISMA rules were followed. The search was conducted from February to April 2026.

Results and Discussion

Summary of Findings

A total of 64 articles were found involving the MeSH Terms. Initially, the exclusion criteria were applied to exclude existing titles and duplications in accordance with the interest described in this work. After this process, the summaries were evaluated, and a new exclusion criterion was applied. A total of 35 articles were evaluated in full, and 26 were included and discussed in this study.

Major Outcomes

Osteoporosis, Osseointegration and Dental Implants - Concept and Epidemiology

Osteoporosis is defined as a systemic skeletal disorder associated with aging, characterized by loss of bone mass, which makes the bone more fragile and more prone to fractures [13-16]. The World Health Organization defined osteoporosis as a level of bone mineral density greater than 2.5 standard deviations below the average of normal young women [17-20]. After 60 years of age, a third of the population has this disorder; it occurs twice as often in women as in men, and its diagnosis is made with greater prevalence from the third decade of life.

Among the systemic alterations, osteoporosis is one of the dysfunctions commonly found by implant dentists [17]. Osteoporosis acts by modifying the metabolism of the bone tissues, disorganizing the trabecular architecture of the cortical and alveolar bone, which are responsible for tooth support. It is estimated that 1.3 million of all fractures and 133,000 of hip fractures occur each year as a result of osteoporosis [20].

Osteoporosis can be classified as type I and type

II. Type I (postmenopausal) occurs when there is loss of trabecular bone mass, resulting in fractures of the vertebrae and wrists, which may be more evident in the mandible and in the alveolar bone, is associated with the aging and plasma decrease of estrogen in the menopause, affecting mainly women; And Type II (senile), occurs when there is loss of trabecular bone mass that can affect both cortical and spongy bone, resulting in hip fractures, which can affect both sexes and in ages over 70 years [12-17].

There is a higher prevalence of the development of osteoporosis in women, and there are some risk factors that may explain this difference, such as early menopause, artificial menopause, nulliparity, and estrogen replacement [21-24]. For men, reduced testicular function (male hypogonadism) can be cited as a risk factor. There are several other risk factors that may predispose both sexes: heredity, tobacco, alcohol, caffeine, obesity, absence of physical activity, ethnicity, changes in calcium levels, malnutrition, decreased levels of vitamin D, elevated Levels of parathyroid hormone and other hormones; all these factors may manifest in both men and women with osteoporosis [13,17].

The recommended intake of calcium is 800 mg day⁻¹, in women who have already gone through menopause, 1.5 g may be required to maintain a positive calcium balance [25,26]. For patients with established osteoporosis, there are drugs that, in general, act directly in the process of bone remodeling, seeking to reduce bone resorption; among them, BP, which are drugs of proven efficacy that act in the prevention and treatment of several Bone diseases [26].

In this sense, dental implants are defined as supports or structures of titanium metal, which, through surgery, are fixed in the maxillary bone, replacing the dental roots, thus allowing the artificial teeth to fit the metal. Dentistry uses several rehabilitation techniques for masticatory functions, and osseointegrated implants are considered safe, provided they are implanted in areas of good quantity and bone quality [12]. However, some systemic conditions may interfere with implant stability, such as osteoporosis. Implantology has shown increasing success rates when it presents a harmonious bone / implant relationship (osseointegration) [12].

The discovery of osseointegration occurred through studies of microcirculation in the bone marrow performed on the rabbit fibula, developed by Per-Ingvar Branemark. He verified in Branemark's studies that a titanium implant, when inserted into the medullary space under certain conditions and remained immobile without mechanical trauma during the period of bone repair, ends up full of compact bone without the interference of other tissues [13-15].

In this context, osteoporosis is a factor that retards

the regeneration of maxillary bone in patients who have undergone implant surgery, prolonging the normal recovery time of maxillary bone that can vary from three to six months [25]. Therefore, it is necessary that people affected by this disease and who will receive dental implants need a longer time for bone repair [25]. Due to the increase in life expectancy, rehabilitation with implants in people over 60 years old is the most common age group in which there is a higher probability of metabolic pathologies [26].

In order to obtain osseointegration of the implant, which is the direct and structural union of the bone tissue to the titanium and function, it is necessary to respect several principles, among them, those related to the surgical technique, respecting tissue physiology [24]. Thus, it is necessary to control the traumatogenic factors during surgery, such as intensity, frequency, and duration of the milling (osteotomies), which can generate excessive trauma to the bone tissue, impairing the bone repair potential of the injured area. Facing situations where the traumatic stimulus exceeds its physiological limit, the implant may be involved by fibrous connective tissues, leading to the formation of a bone or fibrous perimplant interface, without osseointegration [24].

For the success of osseointegratable implants, other factors must also be considered, not only related to the professional (surgical technique), but also the industry and the patient himself. In addition to performing the appropriate surgical technique, it is up to the professional to select the patient, evaluating it as a whole, from his complaint, including his expectation regarding the treatment, mainly comprising his pre-operative systemic and local conditions [25,26]. At the moment of preparation of the receptor bone bed for the subsequent installation of the osseointegrated implant, bone necrosis occurs, which will be replaced by new bone tissue. When there is osteoporosis, the process of bone remodeling can be compromised, preventing or delaying osseointegration [26].

Several authors, including Ourique et al. [17], have already reported on the importance of knowledge of systemic alterations, so that necessary measures are taken to minimize or prevent eventual damage caused by osteoporosis in the anatomical, physiological, and functional integrity of the alveolar bone. All care is necessary for the success of this process, since the immediate benefit of the rehabilitative treatment with implants is observed in the improvement of the capacity to crush food and in the physical and psychological well-being of the patient.

Ishii et al. [12] state that although osteoporosis is a significant factor that can interfere with bone volume and density, it cannot be considered an absolute

contraindication for implant installation. It is essential that during the anamnesis, all patients are questioned about their state of health, reporting the use of medications and the type of medical treatment they are undertaking, so that a safe and effective treatment plan is drawn up for each individual case.

Bisphosphonates (BPs)- Main Considerations

BPs are a widely used drug group for various bone disorders and have been approved by the U.S. Food and Drug Administration for the treatment of osteoporosis, metastatic bone cancer, and Paget's disease [24]. They were first used for industrial purposes in the 19th century to prevent corrosion in the textile, fertilizer, and oil industries. In 1968, the first paper describing the use of BP in medicine was published; however, in 2002, serious side effects of these medications were reported following dental surgery procedures. This includes osteonecrosis, avascular necrosis, osteomyelitis, osteochemionecrosis, and maxillary Bisphosphonate-related osteonecrosis [24].

At the moment, there are two main types of BP: those containing nitrogen (oral: alendronate and risedronate, intravenous: pamidronate and zoledronate) and those that do not contain nitrogen (etidronate, clodronate, and tiludronate). BPs act by suppressing and reducing bone resorption by osteoclasts, directly preventing the recruitment and function of osteoclasts, and indirectly stimulating osteoblasts to produce inhibitors of osteoclast formation [25].

BPs are drugs derived from inorganic pyrophosphate, which is present in the body and physiologically regulates calcification and bone resorption. Pyrophosphate also provides greater resistance to chemical and enzymatic hydrolysis [15]. Camargo, Minosso, and Lopes (2007) [6] report that therapeutic treatment should always combine an anti-resorptive agent with a non-pharmacological measure such as physical exercise and consumption of calcium and vitamin D by diet. Antireabsorption agents are described by Ishii (2009) [12] as estrogen replacement therapy, selective estrogen receptor modulators, BP and calcitonin, and also describes bone formation stimulating agents such as parathyroid hormone.

Also, Ourique et al. [17] have shown in their studies that calcium intake is associated with hormone replacement (estrogen), which leads to an increase in trabecular bone mass. Calcium, when ingested alone, is not able to definitively prevent the onset of osteoporosis. The authors also report that in addition to osteoporosis, age, sex, race, hormonal pattern, decreased vitamin D synthesis, inhibition of calcium absorption, parathormone increase, nicotine, fragile physical structure, renal deficiency, menopause, alcohol,

and low Consumption of calcium may jeopardize the success of an implant.

According to Ishii et al. [12], BPs are anti-resorptive agents derived from pyrophosphonic acid that inhibit bone resorption. Ferreira Junior et al. [10] stated that BP can prevent bone loss, increase bone density, and reduce the risk of fractures resulting from progressive loss of bone mass. In the BP group, alendronate is the most potent because it has an affinity for bone tissue. Another indication to prevent osteoporosis is calcitonin, which is a peptide derived from parafollicular thyroid cells, aiding bone resistance.

Alendronate, for osteoporotic patients, can be administered orally at 10.0 mg / day or 70.0 mg / weekly, and cannot be exceeded because it causes gastrointestinal changes such as erosive esophagitis. It is necessary to use this medicine in fasting, because it is poorly absorbed in the intestine, and to wait 40 to 60 minutes before feeding. It is a drug that deposits about 40-60% rapidly into the bone, and the rest is released through the urine. The plasma half-life of BP is very short, ranging from thirty minutes to two hours, so after these medications are absorbed by the bone tissue, they may persist for more than 10 years in skeletal tissues [17].

A review study with meta-analysis included clinical human studies, randomized or not. A total of 18 publications were included in the review. Regarding implant failure, the meta-analysis found a risk ratio of 1.73 (95% confidence interval [CI] 1.21-2.48, $p = 0.003$) for BP patients when compared to patients who did not take the medicine. The probability of an implant failure in patients receiving BP was estimated at 1.5% (0.015, 95% CI 0.006-0.023, standard error [SE] 0.004, $p < 0.001$). BP cannot be suggested to affect marginal bone loss from dental implants due to a limited number of studies reporting this result. Due to a lack of sufficient information, the meta-analysis for the outcome "postoperative infection" was not performed. The results of the present study cannot suggest that dental implant insertion in patients taking BP affects implant failure rates due to a limited number of published studies, all characterized by a low level of specificity, and most of them dealing with a limited number of cases without an adequate control group. Therefore, the real effect of BP on osseointegration and survival of dental implants is not yet well established [3].

Main Complications - Bisphosphonates

Ishii et al. [12] state that patients who use BP may have impaired healing of the damaged dental implant, as it impedes bone remodeling and may lead to a condition called osteonecrosis, which is considered a side effect of

this drug. Although there are many data on the beneficial effects of BP in the treatment of advanced osseous diseases, numerous reports have documented the ability of these medications to cause local lesions of bone osteonecrosis mainly in the jaw [25].

Osteonecrosis may remain asymptomatic for weeks and possibly months, and lesions usually develop around tapered areas and prior surgical sites, including extractions, retrograde apical tetanus, periodontal surgery, and dental implant surgery. Symptoms include pain, soft tissue edema, infection, tooth loss, and drainage. Radiographically, osteolytic changes are observed, and tissue biopsy shows the presence of actinomycetes [26].

In the dental office, the most common BPs that the implants are exposed to are the oral ones that contain nitrogen, such as risedronate, ibandronate, and alendronate. Comprehensive anamnesis is essential prior to the initiation of any elective treatment; the risk versus benefits of dental treatment should be discussed in detail with the patient [24].

In this context, another study using the BP analyzed the factors related to obtaining effective mechanical and immunological adhesion; viability, epidermal collagen growth factor, and immunoglobulin synthesis were evaluated. The presence of BP culminated in lower cell adhesion to titanium discs, particularly for sodium alendronate (SA) at 5 μ M (40%) and zoledronic acid (ZA) at all concentrations (30 to 50% according to increased concentrations). Reduced cell viability occurred after exposure of these cells to ZA (40%); however, only 5 μ M of SA-treated cells had decreased viability (30%). Reduced synthesis of growth factors and collagen was observed when cells were treated with ZA (20 and 40%, respectively), while about 70% of IgG synthesis was increased. BPs negatively affected adhesion and metabolism of oral mucosal cells, and this effect was related to BP type as well as concentration and treatment period. The negative effects of BPs on oral mucosa cells may hinder the formation of an effective biological seal in osseointegrated implants [4].

Also, a review study aimed to examine the purpose of dental implant placement in patients who have been treated or are undergoing treatment with BP medication. Outcome measures included implant failure or implant-related jaw osteonecrosis. In total, 32 literature sources were reviewed, and 9 of the most relevant articles that fit the criteria were selected. Heterogeneity between studies was found, and no meta-analysis could be performed. Five studies looked at intra-oral BP medication for implant placement, three studies looked at intravenous BP medication for implant placement, and one study evaluated the two types of

medication administered for implant placement. Patients with intraoral therapy appeared to have a better implant survival (5 implants failed 423) rate of 98.8% versus intravenously treated patients (6 implants failed 68) of 91%. The control group, compared with the intraoral

BP group, appeared to have a 97% success rate in implant survival (27 implants failed in 842), showing no significant difference in implant placement success. Patients treated with intravenous BP appear to have a greater chance of developing implant-related jaw osteonecrosis. The intraorally treated group of patients appeared to have more successful results. Implant placement in intraorally treated patients can be considered safe with precautions [5].

Conclusion

In the present research, osteoporosis is a metabolic condition that affects alveolar bone density, but does not present problems for the installation of osseointegrable implants, as long as there is sufficient bone mass in the region where the tooth will be implanted. It suggests to dental surgeons the knowledge of the diagnosis so that it makes a careful evaluation, directing the professional to observe the quality of the bone through routine image examinations. Further, alendronate sodium is used to decrease bone resorption; the drug should be considered as an adjunctive therapeutic agent for the treatment of osteoporosis.

CRedit

Author contributions: **Conceptualization-** All authors; **Investigation-** All authors. **Methodology-** All authors; **Project administration-** All authors; **Supervision-** José Augusto Parola da Cruz; **Writing - original draft-** All authors; **Writing-review & editing-** All authors.

Acknowledgment

Not applicable.

Ethical Approval

Not applicable.

Informed Consent

Not applicable.

Funding

Not applicable.

Data Sharing Statement

No additional data are available.

Publication Statement

All authors have agreed to the publication of this article.

Conflict of Interest

The authors declare no conflict of interest.

Similarity Check

It was applied by Ithenticate@.

Application of Artificial Intelligence (AI)

Not applicable.

Peer Review Process

It was performed.

About The License©

The author(s) 2026. The text of this article is open access and licensed under a Creative Commons Attribution 4.0 International License.

References

1. de Lima Chaparin R, de Lima Medeiros Y, Figueiredo R, Jaguar GC, Lopes RN, Rocha AC, Alves FA. Medication-Related osteonecrosis of the jaw involving dental implants: A 16-Year observational study in a single cancer center. *Clin Oral Investig*. 2025 Sep 2;29(9):436. doi: 10.1007/s00784-025-06519-7.
2. Masri D, Bar-Hai D, Ghanaïem O, Chaushu G, Chaushu L. Peri-Implant-Medication-Related Osteonecrosis of the Jaws: Epidemiological, Clinical and Radiographic Features-A Retrospective Cohort Study. *J Clin Periodontol*. 2026 Jun;53(6):874-886. doi: 10.1111/jcpe.70124.
3. Chrcanovic BR, Albrektsson T, Wennerberg A. Bisphosphonates and dental implants: A meta-analysis. *Quintessence Int*. 2016 Apr;47(4):329-42. doi: 10.3290/j.qi.a35523.
4. Basso FG, Pansani TN, Soares DG, Cardoso LM, Hebling J, de Souza Costa CA. Influence of bisphosphonates on the adherence and metabolism of epithelial cells and gingival fibroblasts to titanium surfaces. *Clin Oral Investig*. 2018 Mar;22(2):893-900. doi: 10.1007/s00784-017-2167-2. Epub 2017 Jul 8.
5. Gelazius R, Poskevicius L, Sakavicius D, Grimuta V, Juodzbals G. Dental Implant Placement in Patients on Bisphosphonate Therapy: a Systematic Review. *J Oral Maxillofac Res*. 2018 Sep 30;9(3):e2. doi: 10.5037/jomr.2018.9302. eCollection 2018 Jul-Sep.
6. Camargo EP, Minosso M, Lopes LC. Caracterização do Uso de Alendronato de Sódio no Tratamento de Osteoporose por Clínicos da Rede Privada de Duas Cidades do Interior de São Paulo. *Rev. Ciênc. Farm. Básica Apl.*, 2007, v.28, n.1, p.77-83.
7. Carvalho PSP. et al. Principais Aspectos da Cirurgia Bucomaxilofacial no Paciente sob Terapia com Bifosfonatos. *RFO UPF* [online]. 2010, vol.15, n.2, pp. 183-189. ISSN 1413-4012.
8. Duarte PM, Nociti Júnior FH. Impacto da Deficiência de Estrógeno e suas Terapias sobre o Tecido Ósseo ao Redor de Implantes de Titânio e na Periodontite Induzida em Ratas Ovariectomizadas. Tese (Doutorado) – Universidade Estadual de Campinas, Faculdade de Odontologia de Piracicaba. Piracicaba, 2004. XVII, 131p.
9. Embracher Filho A. Projeto Colosso: Desenvolvimento de um sistema de implante osseointegrável. Da teoria a prática. Tese (Doutorado) - Faculdade de Odontologia de Araçatuba, UNESP, 2003.
10. Ferreira Júnior CD, Casado PL, Barboza ESP. Osteonecrose Associada aos Bisfosfonatos na Odontologia. *R. Periodontia*. 2007, v.1, n.4.
11. Gegler A. et al. Bifosfonatos e Osteonecrose Maxilar: Revisão da Literatura e Relato de Dois Casos. *Rev. Brasileira de Cancerologia* 2006; 52(1):25-31.
12. Ishii JH. Osteoporose e os Implantes Dentários. São Paulo, 2009. 24 p.
13. Luize DS. et al. A Influência da Osteoporose na Implantodontia. *Arquivos em Odontologia*, Belo Horizonte, 2005, v.41, n.2, p.105-192.
14. Martins MAT et al. Osteonecrose dos Maxilares Associada ao Uso de Bifosfonatos: Importante Complicação do Tratamento Oncológico. *Rev. Bras. Hematol Hemoter* 2009; 31(1):41-46.
15. Migliorati CA et al. O Tratamento de Pacientes com Osteonecrose Associada aos Bifosfonatos. Uma tomada de posição da Academia Americana de Medicina Oral. *J American Dental Association* 2006; 136(12).
16. Misch C E. Implantes Dentais Contemporâneos. Rio de Janeiro: Elsevier. 2008, 3nd.
17. Ourique S.A.M., Ito A.Y., Suarez O.F. Osteoporose em Implantodontia: O Estado Atual da Questão. *Rev. Bras. Implantodontia e Prótese sobre Implantes*, 2005: 12(47/48): 237-45.
18. Goiato MC, Santos DM, Rondon BCS, Moreno A, Baptista GT, Verri FR et al. Care Required When Using Bisphosphonates in Dental Surgical Practice. *J. craniofac. surg*. 2010; 21(6):1966-70.

19. Chadha GK, Ahmadiéh A, Kumar S, Sedghizaded PP. Osseointegration of dental osteonecrosis of the jaw in patients treated with bisphosphonates therapy: a systematic review. *J. oral implantol.* 2013;39(4):510-20.
20. Mellado-Valero A, Ferrer-García JC, CalvoCatalá J, Labaig- Rueda C. Implant treatment in patients with osteoporosis. *Med. oral patol. oral cir. bucal.* 2010; 15:52-7.
21. López-Cedrún JL, Sanromán JF, García A, Peñarrocha M, Feijoo JF, Limeres J, Diz P. Oral bisphosphonate-related osteonecrosis of the jaws in dental implant patients: a case series. *Br. j. oral maxillofac. surg.* 2012;51(8):874-9.
22. Kwon T-G, Lee C-O, Park J-W, Choi S-Y, Rijal G, Shin H-I. Osteonecrosis associated with dental implants in patients undergoing bisphosphonate treatment. *Clin. oral implants res.* 2012; 00:1- 9.
23. Yip JK, Borrell LN, Cho SC, Francisco H, Tarnow DP. Association between oral bisphosphonate use and dental implant failure among middleaged women. *J. clin. periodontol.* 2012;39:408- 14.
24. Memon S, Weltman RL, Katancik JA. Oral Bisphosphonates: Early Endosseous Dental Implant Success and Crestal Bone Changes. A Retrospective Study. *Int. j. oral maxillofac. implants.* 2012;279(5):1216-22.
25. Abtahi J, Tengvall P, Aspenberg P. A bisphosphonate-coating improves the fixation of metal implants in human bone. A randomized trial of dental implants. *Bone.* 2012;50(5):1148-51.
26. Jacobsen C, Metzler P, Rossle M, Obwegeser J, Zemmann W, Gratz KW. Osteopathology induced by bisphosphonates and dental implants: clinical observations. *Clin. oral investig.* 2013;17:167-75.